

Name: _____

Entry number: _____

There are 4 questions for a total of 15 points.

1. Answer the following questions on Propositional Logic.

(a) ($\frac{1}{2}$ point) Fill the truth-table below:

P	Q	R	$P \leftrightarrow Q$	$Q \vee \neg R$	$(P \leftrightarrow Q) \rightarrow (Q \vee \neg R)$
T	T	T	T	T	T
T	T	F	T	T	T
T	F	T	F	F	T
F	T	T	F	T	T
T	F	F	F	T	T
F	T	F	F	T	T
F	F	T	T	F	F
F	F	F	T	T	T

(b) (1 point) Consider the following two compound proposition P, Q :

$$P : (A \vee B) \rightarrow C \quad \text{and} \quad Q : (\neg C \rightarrow \neg A) \vee (\neg C \rightarrow \neg B)$$

Which of the following describe the relationship between P and Q ? Circle all the correct choices and show your reasoning in the space below.

- (a) P and Q are equivalent
 (b) $P \rightarrow Q$
 (c) $Q \rightarrow P$

Solution: We solve this using a truth table.

A	B	C	P	Q
F	F	F	T	T
F	F	T	T	T
F	T	F	F	T
T	F	F	F	T
T	T	F	F	F
T	F	T	T	T
F	T	T	T	T
T	T	T	T	T

- P and Q are not equivalent since the columns for P and Q do not match.
 - $Q \rightarrow P$ does not hold since in the third row, Q evaluates to T but P evaluates to F .
 - $P \rightarrow Q$ holds since there is no row in which P is T but Q is F .
- So, the correct answer is option (b).

2. Answer the following questions on Predicate Logic.

(a) (4 1/2 points) Consider the following predicates:

1. $B(x)$: x is brilliant.
2. $S(x)$: x studies hard.
3. $L(x)$: x is lucky.
4. $C(x, y)$: x clears the final exam of course y .
5. $G(x, y)$: x gets an A grade in course y .
6. $J(x)$: x sleeps too much.

Express each of the statements using quantifiers and the predicates given above. The domain of variable x in the above predicates is the set of all students of COL202 and domain of variable y is the set of all courses being taught at IIT Delhi during Semester-I-2018-19.

	Statement	Quantified expression
S_1	Everyone who clears any final exam studies hard or is brilliant or is lucky.	$\forall x [\exists y C(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))]$
S_2	Everyone who gets an A in some course has cleared the final exam of some course.	$\forall x [\exists y G(x, y) \rightarrow \exists z C(x, z)]$
S_3	No one is lucky.	$\forall x [\neg L(x)]$
S_4	Anyone who sleeps too much does not study hard.	$\forall x [J(x) \rightarrow \neg S(x)]$
S_5	If everyone gets an A in some course, then everyone who sleeps too much is brilliant.	$(\forall x \exists y G(x, y)) \rightarrow (\forall p (J(p) \rightarrow B(p)))$

(b) (1 point) Consider the quantified expressions S_1, \dots, S_5 obtained in the previous part. Use the expressions obtained in the previous part to replace S_1, \dots, S_5 below and then determine whether it makes a valid argument form. Answer “yes” or “no”. You do not need to give explanation for this problem.

$$\begin{array}{l} S_1 \\ S_2 \\ S_3 \\ S_4 \\ \hline \therefore S_5 \end{array}$$

(b) True

Reason (You were not supposed to give this): We obtain the following argument form by replacing the S_1, \dots, S_5 above.

$$\begin{array}{l} \forall x [\exists y C(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))] \\ \forall x [\exists y G(x, y) \rightarrow \exists z C(x, z)] \\ \forall x [\neg L(x)] \\ \forall x [J(x) \rightarrow \neg S(x)] \\ \hline \therefore (\forall x \exists y G(x, y)) \rightarrow (\forall w (J(w) \rightarrow B(w))) \end{array}$$

We will show that the above argument form is valid using rules of inference:

1. $\forall x [\exists y C(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))]$ (Premise)
2. $\forall x [\exists y G(x, y) \rightarrow \exists z C(x, z)]$ (Premise)
3. $\forall x [\neg L(x)]$ (Premise)
4. $\forall x [J(x) \rightarrow \neg S(x)]$ (Premise)
5. $\exists y C(s, y) \rightarrow (S(s) \vee B(s) \vee L(s))$ for an arbitrary student s
(From (1) using Universal instantiation)
6. $\exists y G(s, y) \rightarrow \exists z C(s, z)$
(From (2) using Universal instantiation)
7. $\exists y G(s, y) \rightarrow (S(s) \vee B(s) \vee L(s))$
(From (5) and (6) using modus ponens)
8. $\forall x [\exists y G(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))]$
(From (7) using Universal generalization)
9. $\forall x [(\forall y \neg G(x, y)) \vee S(x) \vee B(x) \vee L(x)]$
(From (8) using De Morgan's law for quantifiers and $p \rightarrow q \equiv p \vee q$)
10. $(\forall y \neg G(s, y)) \vee S(s) \vee B(s) \vee L(s)$ for an arbitrary student s
(From (9) using Universal generalization)
11. $\neg L(s)$
(From (3) using Universal generalization)
12. $(\forall y \neg G(s, y)) \vee S(s) \vee B(s)$
(Resolvent of (10) and (11))
13. $\forall x [\neg J(x) \vee \neg S(x)]$
(From (4) using $p \rightarrow q \equiv p \vee q$)
14. $\neg J(s) \vee \neg S(s)$
(From (13) using Universal generalization)
15. $(\forall y \neg G(s, y)) \vee \neg J(s) \vee B(s)$
(Resolvent of (12) and (14))
16. $\exists x [(\forall y \neg G(x, y)) \vee \neg J(s) \vee B(s)]$
(From (15) using existential generalization)
17. $\forall w \exists x [(\forall y \neg G(x, y)) \vee \neg J(w) \vee B(w)]$
(From (16) using universal generalization)
18. $(\exists x \forall y \neg G(x, y)) \vee (\forall w (J(w) \rightarrow B(w)))$
(From (17) using $p \rightarrow q \equiv p \vee q$)
19. $(\forall x \exists y G(x, y)) \rightarrow \forall w (J(w) \rightarrow B(w))$
(From (18) using $p \rightarrow q \equiv p \vee q$ and De Morgan's law for quantifiers)

Note that step (16) is a correct but a bit unconventional application of existential generalization. In general, if we have a statement $P(s) \vee Q(s)$ that holds for arbitrary s in the domain, then $P(s) \vee (\exists x Q(x)) \equiv \exists x [P(s) \vee Q(x)]$ also holds for an arbitrary element s of the domain. This is the fact that we have used here.

- (c) (2 1/2 points) Consider the quantified expressions S_1, \dots, S_4 obtained in part (a). Use the expressions obtained in part (a) to replace S_1, \dots, S_4 below and then determine whether it makes a valid argument form. Explain your answer. (If your answer is “yes”, then you need to show all steps while using rules of inference)

$$\begin{array}{l} S_1 \\ S_2 \\ S_3 \\ S_4 \\ \hline \therefore \forall x [(\exists y G(x, y)) \rightarrow (J(x) \rightarrow B(x))] \end{array}$$

Solution: We obtain the following argument form by replacing the S_1, \dots, S_4 above.

$$\begin{array}{l} \forall x [\exists y C(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))] \\ \forall x [\exists y G(x, y) \rightarrow \exists z C(x, z)] \\ \forall x [\neg L(x)] \\ \forall x [J(x) \rightarrow \neg S(x)] \\ \hline \therefore \forall x [(\exists y G(x, y)) \rightarrow (J(x) \rightarrow B(x))] \end{array}$$

We will show that the above argument form is valid using rules of inference:

1. $\forall x [\exists y C(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))]$ (Premise)
2. $\forall x [\exists y G(x, y) \rightarrow \exists z C(x, z)]$ (Premise)
3. $\forall x [\neg L(x)]$ (Premise)
4. $\forall x [J(x) \rightarrow \neg S(x)]$ (Premise)
5. $\exists y C(s, y) \rightarrow (S(s) \vee B(s) \vee L(s))$ for an arbitrary student s
(From (1) using Universal instantiation)
6. $\exists y G(s, y) \rightarrow \exists z C(s, z)$
(From (2) using Universal instantiation)
7. $\exists y G(s, y) \rightarrow (S(s) \vee B(s) \vee L(s))$
(From (5) and (6) using modus ponens)
8. $\forall x [\exists y G(x, y) \rightarrow (S(x) \vee B(x) \vee L(x))]$
(From (7) using Universal generalization)
9. $\forall x [(\forall y \neg G(x, y)) \vee S(x) \vee B(x) \vee L(x)]$
(From (8) using De Morgan's law for quantifiers and $p \rightarrow q \equiv p \vee q$)
10. $(\forall y \neg G(s, y)) \vee S(s) \vee B(s) \vee L(s)$ for an arbitrary student s
(From (9) using Universal generalization)
11. $\neg L(s)$
(From (3) using Universal generalization)
12. $(\forall y \neg G(s, y)) \vee S(s) \vee B(s)$
(Resolvent of (10) and (11))
13. $\forall x [\neg J(x) \vee \neg S(x)]$
(From (4) using $p \rightarrow q \equiv p \vee q$)
14. $\neg J(s) \vee \neg S(s)$
(From (13) using Universal generalization)

15. $(\forall y \neg G(s, y)) \vee \neg J(s) \vee B(s)$
(Resolvent of (12) and (14))
16. $\forall x [(\forall y \neg G(x, y)) \vee \neg J(x) \vee B(x)]$
(From (15) using Universal generalization)
17. $\forall x [(\exists y G(x, y)) \rightarrow (J(x) \rightarrow B(x))]$
(From (16) using $p \rightarrow q \equiv p \vee \neg q$)

3. (3 points) Prove or disprove: Let $f : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be a bijection and let $h : \mathbb{N} \times \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$ be a function defined as $h(a, b, c) = f(f(a, b), c)$. Then h is a bijection from $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$ to \mathbb{N} .

Solution: We will prove that the given statement holds. To show that h is a bijection, we need to show that h is one-to-one and onto.

Claim 1: h is a one-to-one function.

Proof. From the definition of one-to-one functions, we need to argue that for any inputs $(a, b, c), (a', b', c') \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$, if $h(a, b, c) = h(a', b', c')$, then $a = a', b = b', c = c'$. Indeed, $h(a, b, c) = h(a', b', c')$ implies that $f(f(a, b), c) = f(f(a', b'), c')$. Since f is one-to-one, this implies that $f(a, b) = f(a', b')$ and $c = c'$. Now using the fact that $f(a, b) = f(a', b')$ and that f is one-to-one, we get that $a = a'$ and $b = b'$. So, we get that if $h(a, b, c) = h(a', b', c')$, then $a = a', b = b',$ and $c = c'$. This completes the proof of the claim. \square

Claim 2: h is onto.

Proof. Using the definition of onto functions, we need to argue that for any $r \in \mathbb{N}$, there exists $(a, b, c) \in \mathbb{N} \times \mathbb{N} \times \mathbb{N}$ such that $h(a, b, c) = r$. Note that since f is an onto function, there exists $(r', c') \in \mathbb{N} \times \mathbb{N}$ such that $f(r', c') = r$. Again, using the fact that f is an onto function, there exists $(a', b') \in \mathbb{N} \times \mathbb{N}$ such that $f(a', b') = r'$. This means that $h(a', b', c') = f(f(a', b'), c') = f(r', c') = r$. This completes the proof of the claim. \square

From Claim 1 and Claim 2, we conclude that h is a bijection.

4. (2 $\frac{1}{2}$ points) Recall the definition of the big-O notation given in the lectures:

Let $f(n)$ and $g(n)$ denote functions mapping positive integers to positive real numbers. The function $f(n)$ is said to be $O(g(n))$ if and only if there exists constants $C, n_0 > 0$ such that for all $n \geq n_0$, $f(n) \leq C \cdot g(n)$.

Prove or disprove: For any functions $f : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ and $g : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ if $f(n)$ is $O(g(n))$, then $5^{f(n)}$ is $O(5^{g(n)})$.

Solution: We will disprove the statement. Consider $f(n) = 2n$ and $g(n) = n$. For these functions we can show that $f(n) = O(g(n))$ since for all $n \geq 1$, $f(n) \leq 2 \cdot g(n)$. However, $5^{f(n)} = 5^{2n}$ and $5^{g(n)} = 5^n$. For any constant $c > 0$, if $c < 1$, then $5^{f(n)} > c \cdot 5^{g(n)}$ for all $n > 0$, otherwise we can show that for all $n \geq \lceil \log_5 c \rceil + 1$, $5^{f(n)} > c \cdot 5^{g(n)}$. This is because if $n \geq \lceil \log_5 c \rceil + 1$, then $5^n > c$, which further implies $5^{2n} > c \cdot 5^n$.